



Cytokine-dependent inflammatory response in SARS-CoV-2 infection

Respuesta inflamatoria dependiente de citoquinas en la infección por SARS-CoV-2

Diana Esperanza Monet Álvarez^{1*} <https://orcid.org/0000-0003-0996-874X>

Emmanuel Zayas Fundora² <https://orcid.org/0000-0002-3830-358X>

Ángel Miguel Aguilar González³ <https://orcid.org/0000-0002-8227-363X>

Gino Iglesias Sordo⁴ <https://orcid.org/0000-0002-9065-1426>

¹Facultad de Medicina No. 1. Universidad de Ciencias Médicas de Santiago de Cuba. Santiago de Cuba, Cuba.

²Universidad de Ciencias Médicas de la Habana. Facultad de Ciencias Médicas “Manuel Fajardo”. La Habana, Cuba.

³Facultad de Ciencias Médicas de Sagua la Grande. Universidad de Ciencias Médicas de Villa Clara. Sagua la Grande, Cuba.

⁴Universidad de Ciencias Médicas de la Habana. Facultad de Ciencias Médicas “10 de octubre”. La Habana, Cuba.

*Autor para la correspondencia. Correo electrónico: esperanza71199@icloud.com

Dear Editor,

COVID-19 still represents a challenge, and a periodic update is necessary, mainly on its pathophysiology as an inflammatory-based disease.

Inflammation is the basis of different clinical entities, and the main reason for consultation in Cuba and the world; known term, commonly seen as a symptom or a risk factor, with an erroneous perception that it is an isolated entity.⁽¹⁾ Both the innate and the acquired immune response act in this process with



numerous local and systemic effects, being a response body homeostasis where the functions of almost all organ systems are modified.⁽²⁾

Different molecules favor its development, such as vasoactive substances, chemokines, lipid metabolites, immunoglobulins, as well as the kinin, coagulation and complement systems, but the main molecules are cytokines;⁽³⁾ small peptides that function as intercellular messengers.

A crucial role is then played by anti-inflammatory cytokines, made up of T-helper 2 lymphocytes, producers of interleukins (IL) 10, 13, 24 and 42, responsible for bringing the defensive response to normal.⁽⁴⁾

Knowing the role of cytokines in inflammation is essential to understand the specific expression of inflammatory-based pathologies, which could mean a significant shift in the management of infectious diseases such as COVID-19.

The arrival of the new coronavirus has generated an unbridled use of epidemiological resources to contain it; however, it will be impossible if the biopsychosocial factors and the conditions of the immune response that are triggered in response to the disease are not known.⁽⁵⁾

SARS-CoV-2 is capable of activating the immune system, giving rise to an uncontrolled systemic inflammatory response, due to the production and secretion of pro-inflammatory cytokines and chemokines by immunoefector cells. This "cytokine storm" is key in the pathophysiology of lung damage, hemodynamic instability, multi-organ failure, and the lethality observed in these patients.⁽⁵⁾

Cytokine storm is a term that, although recently in widespread use, lacks a strict definition. It refers to the overproduction of inflammatory substances with a wide range of biological activity from a variety of tissues and cells, as a result of an infectious process and the loss of negative feedback on the immune system.⁽⁶⁾

A cytokine storm is suspected in all patients with symptoms and signs of COVID-19, associated risk factors, and a torpid course of the disease. It should be taken into account that this entity is associated with a marked elevation of IL-6 (poor prognosis factor in adults with COVID-19) and IL-1, hence the treatment should be aimed at reducing the state of hyperinflammation.⁽⁴⁾

The cytokine storm is not a new concept to immunologists and clinicians, even though it has become more widely known in the wake of COVID-19. They occur in other infections, in autoimmune diseases,



and in people with certain "faulty" genes. However, although it is a previously known term, it is still under study, and many of its treatments are still in the experimental phase.

BIBLIOGRAPHIC REFERENCES

1. Porter D. Inflamación. American Academy of Ophthalmology; 2020. [acceso: 24/01/2022]. Disponible en: <https://www.aao.org/salud-ocular/sintomas/inflamacion>
2. MiSistemaInmune. Inflamación: tipos y patologías asociadas. MiSistemaImmune; 2019. [acceso: 24/01/2022]. Disponible en: [https://www.misistemainmune.es/enfermedades-sistema-inmunitario/inflamatorias/inflamacion-tipos-y-patologias-asociadas#:~:text=La%20inflamaci%20n%20es%20una%20respuesta,alguna%20sustancia%20agresiva\)%2C%20etc](https://www.misistemainmune.es/enfermedades-sistema-inmunitario/inflamatorias/inflamacion-tipos-y-patologias-asociadas#:~:text=La%20inflamaci%20n%20es%20una%20respuesta,alguna%20sustancia%20agresiva)%2C%20etc)
3. Sosa-Medellín MÁ, Ponce-Mendoza RA, Luviano-García JA, Almanza-Valdez HE, Maheda-García HJ, Santos-Macías JE, et al. COVID-19: ¿tormenta de citocinas o sepsis viral? Med Int Mex. 2021 [acceso: 24/01/2022]; 37(4):580-5. Disponible en: <https://www.medigraphic.com/pdfs/medintmex/mim-2021/mim214m.pdf>
4. Miranda Pedroso R. Tormenta de citoquinas en la infección por SARS-CoV-2 (COVID19). Revista Cubana de Medicina Intensiva y Emergencias. 2021 [acceso: 24/01/2022]; 20(3):e830 Disponible en: <http://www.revmie.sld.cu/index.php/mie/article/view/830/pdf>
5. Hurtado de Mendoza Amat JD, Montero González T, Borrajero Martínez I, Capo de Paz V, López Marín L, Dominguez Alvarez C. Respuesta inflamatoria sistémica de la COVID-19 y su expresión morfológica, el daño múltiple de órganos. Revista Cubana de Medicina Militar. 2021 [acceso: 02/01/2022]; 50(3): 02101592. Disponible en: <http://www.revmedmilitar.sld.cu/index.php/mil/article/view/1592/1038>
6. Noris-García E, Robinson-Agramonte Md. Psiconeuroinmunoendocrinología y COVID-19. Rev. Electron. Dr. Zoilo Marinello Vidaurreta. 2021 [acceso: 02/01/2022]; 46(1): [aprox. 4 p.]. Disponible en: <http://revzoilomarinello.sld.cu/index.php/zmv/article/view/2523>



Conflict of interest

No conflict of interest is declared and no funding was received for the development of this article.